

8th WCCE, 28 August 2009, Montréal

Transport properties of hydrogen bonding fluids: molecular simulation and Taylor dispersion measurements

Gabriela Guevara-Carrión¹, Jadran Vrabec², Hans Hasse¹

¹ Laboratory of Engineering Thermodynamics (LTD)
University of Kaiserslautern, Germany

² Thermodynamics and Energy Technology (ThET)
University of Paderborn, Germany

Molecular simulation

- **Goal**

Macroscopic behaviour from intermolecular interactions


- **Molecular Dynamics (MD)**

Solve Newton's equation of motion

- ✓ Static properties
- ✓ Dynamic properties

- **Transport properties prediction**

- ✓ Equilibrium MD (EMD)
 - Green-Kubo method
 - D_i , D_{ij} , η



Molecular models

Rigid United-Atom multicenter Lennard-Jones (LJ) models


Hydrogen bond modelling through point charges

Mixtures: unlike LJ parameters

→ Lorentz-Berthelot

$$\sigma_{AB} = \left(\frac{\sigma_A + \sigma_B}{2} \right)$$

$$\epsilon_{AB} = \sqrt{\epsilon_A \epsilon_B}$$



• Parameters

methanol ethanol water

| | | | |
|----------------------------|---|---|--------------------------------------|
| ✓ Geometric parameters | 3 | 5 | SPC/E |
| ✓ Lennard-Jones parameters | 4 | 6 | Berendsen et al., 1987 TIP4P |
| ✓ Point charge parameters | 2 | 2 | Jorgensen et al., 1983 TIP4P_2005 |
| | | | Abascal & Vega, 2005 |

Parameters adjusted to:

- ✓ vapour pressure
- ✓ sat. vapour density
- ✓ critical temperature

**No transport
properties !**

EMD: Green-Kubo formalism

equilibrium
fluctuations



$$F_i = \sum_j L_{ij} Y_j$$

microscopic flux



microscopic
equilibrium

transport coefficients



autocorrelation functions

- self-diffusion

$$D_i = \frac{1}{3N_i} \int_0^{\infty} dt \left\langle \sum_i^N v_i(0) \cdot v_i(t) \right\rangle$$

- binary Maxwell-Stefan mutual diffusion





$$D_{12} = \frac{x_2}{3N_1} \left(\frac{M_1 x_1 + M_2 x_2}{M_2 x_2} \right)^2 \int_0^{\infty} dt \left\langle \sum_{i=1}^{N_1} v_i(0) \cdot \sum_{j=1}^{N_1} v_j(t) \right\rangle$$

- shear viscosity



$$\eta_s = \frac{1}{Vk_B T} \int_0^{\infty} dt \left\langle J_p^{xy}(t) \cdot J_p^{xy}(0) \right\rangle$$

$$J_p^{xy} = \sum_{i=1}^N m_i \cdot v_i^x \cdot v_i^y - \sum_{i=1}^N \sum_{j>i}^N r_{ij}^x \frac{\partial \phi(r_{ij})}{\partial r_{ij}^y}$$

Autocorrelation functions





Prediction transport coefficients: methanol (I)





● Prediction MD simulation
 + Experiments (literature)
 all data at 0.1 MPa

Prediction transport coefficients: methanol (II)





- Prediction MD simulation
 - + Experiments (literature)
- all data at 100 MPa**

Prediction transport coefficients: ethanol




● Prediction MD simulation
 + Experiments (literature)
 all data at 0.1 MPa

Prediction transport coefficients: water




- ▲ Prediction simulation SPC/E model
 - Prediction simulation TIP4P model
 - Prediction simulation TIP4P_2005 model
 - + Experiments (literature)
- all data at 0.1 MPa

Taylor dispersion fundamentals




Concentration distribution

fast diffusion



slow diffusion




$$D_{\text{narrow}} > D_{\text{broad}}$$

Taylor's solution

$$\bar{c}_i(t) = \frac{V_i c_i}{2\pi R^2} \frac{1}{\sqrt{(\pi k t)}} \exp\left(\frac{-L^2 \left(1 - \frac{t}{\tau}\right)^2}{4kt}\right)$$


$$k = \frac{R^2 L^2}{48 \tau^2 D_F}$$

Taylor dispersion equipment




Fickian diffusion coefficients from Taylor dispersion measurements (I)

methanol + water (298 K)




ethanol + water (298 K)




Fickian diffusion coefficient prediction vs. experiment

methanol + water (298K)




ethanol + water (298K)



● Prediction MD simulation
+ Experiments (literature)


all data at 0.1 MPa

Fickian diffusion coefficient from Taylor dispersion measurements (II)




Self-diffusion coefficients in mixtures


methanol + water



ethanol + water



methanol + ethanol




- Prediction MD simulation
- + Experiments (literature)


all data at 298 K and 0.1 MPa

Shear viscosity of mixtures


methanol + water



ethanol + water




ethanol + methanol




- Prediction MD simulation
 - + Experiments (literature)
- all data at 298 K and 0.1 MPa

Self-diffusion coefficient: other substances


ammonia



methylamine




dimethylether




- Prediction MD simulation
 - + Experiments (literature)
- all data at 10 MPa

Shear viscosity: other substances


ammonia



methylamine



dimethylether



- Prediction MD simulation
 - + Experiments (literature)
 - DDTB (correlation)
- all data at 10 MPa

Summary

- Prediction of transport properties of hydrogen-bonding liquids using rigid UA molecular models
- Self-diffusion, mutual diffusion, and shear viscosity
- Green-Kubo MD method
- Pure substances and mixtures
- Very good agreement using models that were adjusted just to VLE data
- Best water model tested from the literature: TIP4P_2005
- Experimental determination of transport diffusion coefficients using the Taylor dispersion method